

Warning Statements

Dust.

This product contains Dust. Do not swallow. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Fertilisers Containing Cadmium, Lead, Mercury.

If this fertiliser contains >1mg/kg cadmium, and/or >20mg/kg lead and/or >0.2mg/kg mercury, the following warning applies.

This product contains heavy metal impurities. Its use may lead to an accumulation of cadmium, and/or lead and/or mercury in the soil. Depending on soil characteristics, irrigation water quality, plant species and variety, crop uptake of cadmium and/or lead and/or mercury may exceed the maximum limits allowed in food standards. In pasture, the offal from grazing animals may also exceed these limits.

Fertilisers Containing >100mg Cd/kg P

This product should not be fed to livestock or used in stock feed mixtures.

Fertilisers Containing Fluorine.

This product contains fluorine as an impurity. Do not feed this product to livestock or use in stock feed mixtures. If top dressing pastures do not graze for 3 weeks or until rain or irrigation is received.

Fertilisers Containing Molybdenum.

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Fertilisers Containing Trace Elements

Misuse of trace elements can cause problems with livestock. It is recommended that when applying trace elements, livestock should be withheld from the treated area until either after an irrigation or until 10mm of rain has fallen or 21 days has elapsed after trace element application. When used at above Impact Fertilisers normal suggested rate, may induce toxic levels in soil, plants or animals. May damage seed or emerging seedlings.

Urea

This fertilizer contains biuret. It is not recommended for repeated foliar applications on annual or perennial horticultural crops.

Blends Containing Urea

Limited compatibility. Blend and use. Do not store. Apply when humidity is low.

Storage:

Do not store product in silos.

Fertilisers Containing Selenium

Excessive use of selenium (Se) can be toxic to livestock. This product should not be used if selenium deficiency does not exist in stock grazing on the area where it is to be applied. Do not allow stock access to spilt or unused fertilizer. This product must not be applied at a rate greater than that stated on the label or advice note.